



Adult female gypsy moth with egg mass

A native of Europe, the gypsy moth (*Lymantria dispar*) was accidentally released in Massachusetts in 1869. Infestations of the pest have gradually spread, leaving behind millions of acres of defoliated trees. Since 1980, the gypsy moth has defoliated more than one million acres in Maryland. During this period, the Gypsy Moth Cooperative Suppression Program sprayed the trees with carefully selected insecticides on another 1.8 million acres statewide. The suppression spray program has protected the trees from severe leaf loss on an average of over 97 percent of the acreage treated each year.

From the early 1980s to the early 1990's, severe infestations of gypsy moth caterpillars and the resultant defoliation occurred primarily in Allegany, Anne Arundel, Baltimore, Carroll, Cecil, Frederick, Garrett, Harford, Howard, Kent, Montgomery, Prince George's, and Washington counties. Most of the Maryland Department of Agriculture's (MDA) gypsy moth suppression activities were conducted in these counties. By 1994, the northern infestations had collapsed, but on the Eastern Shore and in Southern Maryland, the caterpillars were very active and the suppression spraying was conducted largely in those areas. Although gypsy moth caterpillar populations were low between 1996 and 1999, MDA's annual fall survey program detected several increasing populations throughout the state.

Gypsy moth caterpillar populations rebounded significantly in the spring of 2000 – defoliating 22,824 acres – and again in 2001 – defoliating 46,183 acres. In 2000, the MDA's Gypsy Moth Suppression Program sprayed the trees on 16,971 acres. In 2001, the Suppression Program sprayed the trees on 48,588 acres.

In 2002, MDA sprayed 39,134 acres of trees statewide and only 112 acres of trees (untreated) were defoliated. As populations again began to collapse, spray acreage was reduced to 14,053 acres in 2003 and to 660 acres in 2004. There was no suppression spraying in 2005.

In 2006 and 2007, conditions seem to have been especially favorable for gypsy moth larvae. The large, healthy caterpillar populations fed voraciously on the oaks and other hardwoods in Maryland, defoliating the trees on 15,793 acres in the spring of 2006 and on 68,460 acres in the spring of 2007.

In 2006, in response to population data gathered in the fall of 2005, the MDA's Gypsy Moth Suppression Program sprayed the trees on 25,456 acres statewide. In 2007, the Suppression Program sprayed the trees on 50,173 acres statewide.

Responding to the defoliation in 2007 and to population predictions from the 2007 fall egg mass survey program, MDA's Gypsy Moth Suppression Program sprayed 99,222 acres in the spring of 2008. Statewide that year, the caterpillars defoliated the trees on 19,279 acres. In the ensuing years the population has continued to cycle up and down. The year 2009 brought a need to treat 32,543 acres, while in 2010 only 144 acres were treated, and no areas needed treatment in 2011. Acres treated rebounded to 2,303 in 2012, increased to 11,996 acres in 2013, and 5,164 acres in 2014. Gypsy moth populations go through cycles, but are heavily influenced by seasonal weather conditions, and can fluctuate from a low to a high potentially damaging level in as little as two years.



Suburban yard in late June - trees were defoliated by gypsy moth caterpillars.



Gypsy moth egg mass


Gypsy moth caterpillars do all the harm. They hatch in large numbers - an egg mass can contain more than 1000 eggs - and the caterpillars feed ravenously on leaves.

Additionally, because they are not native to North America, they have few natural enemies and feed and reproduce in relative safety. Where populations are high and no control measures are taken, trees can be heavily defoliated.


Trees that lose more than 60 percent of their leaves often re-leaf depleting their energy reserves and leaving them under severe stress and vulnerable to attack by diseases and damaging insects. The combination of defoliation, re-leafing, and disease or insect activity, complicated by other factors such as drought or soil compaction, may kill the tree in one to three years. To protect valuable forest and shade trees, the Maryland Department of Agriculture maintains a monitoring and control program to manage the pest. Each fall MDA personnel survey all qualifying areas for gypsy moth egg masses to determine where populations of the pest might be growing. Natural enemies have been introduced statewide, but the most effective suppression results from spraying trees in infested areas with carefully selected insecticides to destroy the caterpillars early in their development.

In the more than 100 years since the gypsy moth was introduced into North America, it has steadily spread in spite of man's best efforts to control it. No one agency can hope to eliminate it, but by combining efforts and employing a variety of controls, property owners, neighborhood organizations, businesses, and government agencies can hope to limit its effects.

**Trees preferred by gypsy moth**

- All oaks
  - Apple
  - American Beech
  - Birch
  - Sweetgum
  - Linden
  - Willow
  - Hawthorn
- 

**Trees not preferred by gypsy moth**

- American
  - Catalpa
  - Red Spruce
  - Sycamore
  - Holly
  - Tulip Poplar
  - Locust
- 

**Maryland Department of Agriculture  
Forest Pest Management Regional Offices**

**Northeast Office**

P.O. Box 502  
Forest Hill, Maryland 21050  
410-879-8034  
northeastfpm@gmail.com  
Baltimore City, Baltimore, Cecil, Harford,  
Kent counties

**Eastern Shore Office**

28577 Mary's Ct., Suite 4  
Easton, MD 21601  
heather.disque@maryland.gov  
410-822-8234  
Caroline, Dorchester, Queen Anne's, Talbot,  
Wicomico, Worcester counties

**Central Office**

92 Thomas Johnson Dr., Suite 110  
Frederick, Maryland 21702  
240-529-0390  
mdacent@erols.com  
Carroll, Frederick, Howard, Montgomery,  
Eastern Washington counties

**Southern Office**

50 Harry S. Truman Parkway  
Annapolis, MD 21401  
fpm.mda@maryland.gov  
410-841-5922  
Anne Arundel, Calvert, Charles,  
Prince George's, St. Mary's counties

**Western Office**

3 Pershing Street, Room 100  
Cumberland, Maryland 21502  
301-777-3601  
mdafpmwestern@gmail.com  
Allegany, Garrett, Western Washington counties

**GYPSY MOTH  
IN  
MARYLAND**



**Late instar gypsy moth caterpillar.**

**Maryland Department of Agriculture  
Forest Pest Management Section  
50 Harry S. Truman Parkway  
Annapolis, Maryland 21401  
www.mda.maryland.gov**



**Larry Hogan, Governor  
Boyd Rutherford, Lt. Governor  
Joseph Bartenfelder, Secretary  
James P. Eichhorst, Deputy Secretary**

Note: there are many trees, such as maple, elm, cherry, hickory, blue spruce and white pine, as well as several shrubs and ornamental plants that are less desirable to gypsy moth but are still attacked and often defoliated.

